

Trifolium squamosum

Habitus and growth type

Plant height [m]: **0.29**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **1.2**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8.6**

Reaction indicator value: **6.2**

Salinity indicator value: **1.5**

Disturbance indicator values

Disturbance frequency: **1.01**

Disturbance frequency (herb layer): **2.04**

Disturbance severity: **0.55**

Disturbance severity (herb layer): **0.5**

Mowing frequency: **0.45**

Grazing pressure: **0.39**

Soil disturbance: **0.2**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [MF \(JUN\) Juncetea maritimi](#)

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Grassland (non-alpine, non-saline), Scrub**