

# *Trichophorum pumilum*

## Habitus and growth type

Plant height [m]: **0.09**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

## Flower

Flowering period: **July-August**

## Fruit, seed and dispersal

Seed mass [mg]: **0.56**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **2**

Moisture indicator value: **8.3**

Nutrient indicator value: **1.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **2.02**

Disturbance frequency (herb layer): **2.02**

Disturbance severity: **0.21**

Disturbance severity (herb layer): **0.21**

Mowing frequency: **0.6**

Grazing pressure: **0.13**

Soil disturbance: **0.01**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [PA \(SCH\) \*Scheuchzerio palustris-Caricetea fuscae\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [Q46 Carpathian travertine fen with halophytes](#)

Constant species of EUNIS habitats: [Q46 Carpathian travertine fen with halophytes](#)

Dominant species of EUNIS habitats: [Q46 Carpathian travertine fen with halophytes](#)

### Broad habitat

Occurrence in broad habitats: **Spring, Mire, Sparsely vegetated (incl. rock and scree)**