

# ***Tripidium ravennae***

## **Habitus and growth type**

Plant height [m]: **2.5**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## **Leaf**

Specific leaf area [mm<sup>2</sup>/mg]: **6.41**

## **Flower**

Flowering period: **June-October**

## **Fruit, seed and dispersal**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### **Environmental relationships**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

### **Ellenberg-type indicator values**

Light indicator value: **8.3**

Temperature indicator value: **9.3**

Reaction indicator value: **8**

Nutrient indicator value: **6.7**

Salinity indicator value: **2**

### **Disturbance indicator values**

Disturbance frequency: **0.87**

Disturbance frequency (herb layer): **1.83**

Disturbance severity: **0.47**

Disturbance severity (herb layer): **0.36**

Mowing frequency: **0.22**

Grazing pressure: **0.21**

Soil disturbance: **0.19**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FD \(NER\) \*Nerio-Tamaricetea\*](#), [MF \(JUN\) \*Juncetea maritimi\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [N1J Mediterranean and Black Sea moist and wet dune slack](#)

Constant species of EUNIS habitats: [N1J Mediterranean and Black Sea moist and wet dune slack](#)

### Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Wetland, Saline vegetation, Scrub**