

# *Trisetum alpestre*

## Habitus and growth type

Plant height [m]: **0.3**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **June-August**

## Fruit, seed and dispersal

Seed mass [mg]: **0.28**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **2.9**

Moisture indicator value: **4.4**

Reaction indicator value: **8**

Nutrient indicator value: **2.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.07**

Disturbance frequency (herb layer): **1.39**

Disturbance severity: **0.33**

Disturbance severity (herb layer): **0.29**

Mowing frequency: **0.02**

Grazing pressure: **0.25**

Soil disturbance: **0.11**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CU \(SES\) \*Elyno-Seslerietea\*](#), [KC \(ASP\) \*Asplenietea trichomanis\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R44 Arctic-alpine calcareous grassland](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **6**

Continentality amplitude: **2**