

Trisetum spicatum

Habitus and growth type

Plant height [m]: **0.11**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.42**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **1**

Moisture indicator value: **5**

Reaction indicator value: **6.3**

Nutrient indicator value: **1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.6**

Disturbance frequency (herb layer): **1.66**

Disturbance severity: **0.29**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0**

Grazing pressure: **0.2**

Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**KE \(THL\) *Thlaspietea rotundifolii***](#),

[**QH \(ARC\) *Matricario-Poetea arcticae***](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**U21 Boreal and Arctic siliceous scree and block field**](#), [**U25 Boreal and Arctic base-rich scree and block field**](#)

Constant species of EUNIS habitats: [**U21 Boreal and Arctic siliceous scree and block field**](#), [**U25 Boreal and Arctic base-rich scree and block field**](#)

Broad habitat

Occurrence in broad habitats: **Oromediterranean scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**