

Triticum aestivum

Habitus and growth type

Plant height [m]: **0.94**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **22.78**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **41.18**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Archeophyte**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.2**

Temperature indicator value: **7**

Moisture indicator value: **4.7**

Reaction indicator value: **6.7**

Nutrient indicator value: **6.2**

Salinity indicator value: **0**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [V11 Intensive unmixed crops](#), [V13 Arable land with unmixed crops grown by low-intensity agricultural methods](#)

Constant species of EUNIS habitats: [V11 Intensive unmixed crops](#), [V13 Arable land with unmixed crops grown by low-intensity agricultural methods](#)

Dominant species of EUNIS habitats: [V11 Intensive unmixed crops](#), [V13 Arable land with unmixed crops grown by low-intensity agricultural methods](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**