Trochiscanthes nodiflora

Habitus and growth type

Plant height [m]: **1.15**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **4.04**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **4**
Temperature indicator value: **4.3**
Moisture indicator value: **5**
Reaction indicator value: **6**
Nutrient indicator value: **5.5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.02**
Disturbance frequency (herb layer): **1.32**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.11**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0**
Grazing pressure: **0.2**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CA (FAG) *Carpino-Fagetea sylvaticae***

Broad habitat

Occurrence in broad habitats: **Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz