

Vaccinium oxycoccos

Habitus and growth type

Plant height [m]: **0.2**

Life span: **Perennial**

Life form: **Chamaephyte**

Leaf

Leaf type (woody plants only): **Sclerophyllous**

Leaf phenology (woody plants only): **Evergreen**

Specific leaf area [mm²/mg]: **16.24**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **13.48**

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.4**

Temperature indicator value: **4.4**

Moisture indicator value: **8.8**

Reaction indicator value: **2.6**

Nutrient indicator value: **1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.56**

Disturbance frequency (herb layer): **1.33**

Disturbance severity: **0.31**

Disturbance severity (herb layer): **0.11**

Mowing frequency: **0.14**

Grazing pressure: **0.13**

Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [PB \(OXY\) Oxycocco-Sphagnetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [Q11 Raised bog](#), [Q22 Poor fen](#), [Q25 Non-calcareous quaking mire](#), [Q42 Extremely rich moss-sedge fen](#), [T3J Pinus and Larix mire forest](#)

Constant species of EUNIS habitats: [P3g Dystrophic-water vegetation](#), [Q11 Raised bog](#), [Q21 Oceanic valley mire](#), [Q22 Poor fen](#), [Q25 Non-calcareous quaking mire](#), [Q42 Extremely rich moss-sedge fen](#), [Q44 Calcareous quaking mire](#), [T16 Broadleaved mire forest on acid peat](#), [T3J Pinus and Larix mire forest](#), [T3K Picea mire forest](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Mire, Forest**

Distribution

Continentality amplitude: **6**