

Vaccinium uliginosum

Habitus and growth type

Plant height [m]: **0.42**

Life span: **Perennial**

Life form: **Chamaephyte, Dwarf shrub**

Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**

Leaf phenology (woody plants only): **Winter deciduous**

Specific leaf area [mm²/mg]: **14.45**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **121.08**

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.6**

Temperature indicator value: **3.2**

Reaction indicator value: **1.7**

Nutrient indicator value: **2.1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.54**
 Disturbance frequency (herb layer): **1.23**
 Disturbance severity: **0.39**
 Disturbance severity (herb layer): **0.2**
 Mowing frequency: **0.02**
 Grazing pressure: **0.2**
 Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [AC \(LOI\) *Loiseleurio procumbentis-Vaccinietea*](#), [PB \(OXY\) *Oxycocco-Sphagnetea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [Q31 Palsa mire](#), [S11 Shrub tundra](#), [S12 Moss and lichen tundra](#), [S22 Alpine and subalpine ericoid heath](#), [T3J Pinus and Larix mire forest](#)

Constant species of EUNIS habitats: [N1F Baltic coniferous coastal dune forest](#), [Q11 Raised bog](#), [Q23 Relict mire of Mediterranean mountains](#), [Q31 Palsa mire](#), [Q45 Arctic-alpine rich fen](#), [R43 Temperate acidophilous alpine grassland](#), [S11 Shrub tundra](#), [S12 Moss and lichen tundra](#), [S22 Alpine and subalpine ericoid heath](#), [S23 Alpine and subalpine Juniperus scrub](#), [S26 Subalpine Pinus mugo scrub](#), [T16 Broadleaved mire forest on acid peat](#), [T1C Temperate and boreal mountain Betula and Populus tremula forest on mineral soils](#), [T34 Temperate subalpine Larix, Pinus cembra and Pinus uncinata forest](#), [T3G Pinus sylvestris light taiga](#), [T3H Larix light taiga](#), [T3J Pinus and Larix mire forest](#), [T3K Picea mire forest](#), [U25 Boreal and Arctic base-rich scree and block field](#), [U35 Boreal and Arctic base-rich inland cliff](#)

Broad habitat

Occurrence in broad habitats: **Spring, Mire, Alpine-subalpine and arctic grassland, Heathland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**
 Continentality amplitude: **6**