Valeriana stolonifera

Habitus and growth type

Plant height [m]: **0.8**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.1**
Temperature indicator value: **6.3**
Moisture indicator value: **4.3**
Reaction indicator value: **7**
Nutrient indicator value: **4**
Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **0.49**
Disturbance frequency (herb layer): **1.56**
Disturbance severity: **0.54**
Disturbance severity (herb layer): **0.33**
Mowing frequency: **0.31**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.23**
Soil disturbance: **0.12**

Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz