

Valerianella carinata

Habitus and growth type

Plant height [m]: **0.2**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **32.97**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.55**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**

Moisture indicator value: **4.1**

Reaction indicator value: **7.6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.17**

Disturbance frequency (herb layer): **1.96**

Disturbance severity: **0.62**

Disturbance severity (herb layer): **0.57**

Mowing frequency: **0.45**

Grazing pressure: **0.29**

Soil disturbance: **0.42**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CK \(SED\) *Sedo-Scleranthetea*, QC \(CHE\) *Chenopodietea*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **6**