

# *Valerianella coronata*

## Habitus and growth type

Plant height [m]: **0.25**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **47.81**

## Flower

Flowering period: **March-June**

## Fruit, seed and dispersal

Seed mass [mg]: **1.16**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate reaction relationship: **Slightly acidic to near-neutral**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Moisture indicator value: **3.1**

Reaction indicator value: **6.4**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.42**

Disturbance frequency (herb layer): **2.11**

Disturbance severity: **0.7**

Disturbance severity (herb layer): **0.67**

Mowing frequency: **0.49**

Grazing pressure: **0.3**

Soil disturbance: **0.41**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [QC \(CHE\) \*Chenopodietea\*](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**