

# *Veronica anagalloides*

## Habitus and growth type

Plant height [m]: **0.27**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **19.58**

## Flower

Flowering period: **May-October**

## Fruit, seed and dispersal

Seed mass [mg]: **0.03**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.7**

Moisture indicator value: **8.7**

Salinity indicator value: **0.5**

### Disturbance indicator values

Disturbance frequency: **1.7**

Disturbance frequency (herb layer): **1.92**

Disturbance severity: **0.58**

Disturbance severity (herb layer): **0.57**

Mowing frequency: **0.2**

Grazing pressure: **0.23**

Soil disturbance: **0.27**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [OC \(ISO\) Isoëto-Nanojuncetea](#)

Broad habitat

Occurrence in broad habitats: **Spring, Wetland**

## **Distribution**

Continentality: **7**

Continentality amplitude: **6**