

# *Veronica bachofenii*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Disturbance indicator values

Disturbance frequency: **0.5**

Disturbance frequency (herb layer): **1**

Disturbance severity: **0.39**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.11**

Grazing pressure: **0.2**

Soil disturbance: **0.12**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [KC \(ASP\) \*Asplenietea trichomanis\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [U32 Temperate high-mountain siliceous inland cliff](#)

Constant species of EUNIS habitats: [U32 Temperate high-mountain siliceous inland cliff](#)

Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree)**