

# *Veronica scutellata*

## Habitus and growth type

Plant height [m]: **0.21**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **30.24**

## Flower

Flowering period: **May-September**

## Fruit, seed and dispersal

Seed mass [mg]: **0.28**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.2**

Temperature indicator value: **4.8**

Moisture indicator value: **9.1**

Reaction indicator value: **3.9**

Nutrient indicator value: **2.9**

Salinity indicator value: **0.2**

### Disturbance indicator values

Disturbance frequency: **1.44**

Disturbance frequency (herb layer): **1.74**  
Disturbance severity: **0.38**  
Disturbance severity (herb layer): **0.37**  
Mowing frequency: **0.33**  
Grazing pressure: **0.2**  
Soil disturbance: **0.14**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: **[OB \(LIT\) Littorelletea uniflorae](#)**

### Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Wetland, Mire, Grassland (non-alpine, non-saline), Scrub**

## **Distribution**

Continentality amplitude: 7