

# *Vicia barbazitae*

## Habitus and growth type

Plant height [m]: **0.4**

Life span: **Annual, Perennial**

Life form: **Hemicryptophyte, Therophyte**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **4**

Temperature indicator value: **6**

Moisture indicator value: **4**

Reaction indicator value: **5**

Nutrient indicator value: **5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.12**

Disturbance frequency (herb layer): **1.41**

Disturbance severity: **0.72**

Disturbance severity (herb layer): **0.26**

Mowing frequency: **0.09**

Grazing pressure: **0.23**

Soil disturbance: **0.18**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [\*\*CB \(PUB\) Quercetea pubescentis\*\*](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**