Vicia bithynica

Habitus and growth type

Plant height [m]: **0.34**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Seed mass [mg]:
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.2**
Moisture indicator value: **4.7**
Reaction indicator value: **6.5**
Nutrient indicator value: **5.5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.4**
Disturbance frequency (herb layer): **1.56**
Disturbance severity: **0.7**
Disturbance severity (herb layer): **0.46**
Mowing frequency: **0.28**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.23**
Soil disturbance: **0.26**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QC (CHE) *Chenopodietea***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz