

Vicia disperma

Habitus and growth type

Plant height [m]: **0.3**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Seed mass [mg]: **15.1**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **10**

Moisture indicator value: **3**

Reaction indicator value: **5**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.35**

Disturbance frequency (herb layer): **1.49**

Disturbance severity: **0.68**

Disturbance severity (herb layer): **0.36**

Mowing frequency: **0.16**

Grazing pressure: **0.25**

Soil disturbance: **0.21**

Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**