

Vicia narbonensis

Habitus and growth type

Plant height [m]: **0.44**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **245.1**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.4**

Moisture indicator value: **4.6**

Reaction indicator value: **7.3**

Nutrient indicator value: **5.3**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **0.68**

Disturbance frequency (herb layer): **1.6**

Disturbance severity: **0.71**

Disturbance severity (herb layer): **0.56**

Mowing frequency: **0.36**

Grazing pressure: **0.18**

Soil disturbance: **0.4**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**

Distribution

Continentality amplitude: 7