

Vicia pannonica

Habitus and growth type

Plant height [m]: **0.42**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **36.9**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.4**

Moisture indicator value: **3.8**

Reaction indicator value: **6.5**

Nutrient indicator value: **5.2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.42**

Disturbance frequency (herb layer): **1.92**

Disturbance severity: **0.72**

Disturbance severity (herb layer): **0.69**

Mowing frequency: **0.61**

Grazing pressure: **0.2**

Soil disturbance: **0.5**

Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Scrub, Sparsely vegetated (incl. rock and scree),**

Synanthropic