

# *Vicia peregrina*

## Habitus and growth type

Plant height [m]: **0.3**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **21.82**

## Flower

Flowering period: **February-May**

## Fruit, seed and dispersal

Seed mass [mg]: **41.7**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **8**

Moisture indicator value: **3.3**

Reaction indicator value: **7**

Nutrient indicator value: **5.3**

Salinity indicator value: **0.5**

### Disturbance indicator values

Disturbance frequency: **1.35**

Disturbance frequency (herb layer): **2**

Disturbance severity: **0.75**

Disturbance severity (herb layer): **0.73**

Mowing frequency: **0.73**

Grazing pressure: **0.26**

Soil disturbance: **0.47**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [QC \(CHE\) \*Chenopodietea\*](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**