

Vinca minor

Habitus and growth type

Plant height [m]: **0.3**

Life span: **Perennial**

Life form: **Chamaephyte**

Leaf

Specific leaf area [mm²/mg]: **24.7**

Flower

Flowering period: **January-April**

Fruit, seed and dispersal

Seed mass [mg]: **7.38**

Dispersal mode: **Myrmecochory, Anthropochory**

Dispersal distance class: **3, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **3.9**

Temperature indicator value: **5.9**

Moisture indicator value: **5.3**

Reaction indicator value: **6.6**

Nutrient indicator value: **6.2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.04**

Disturbance frequency (herb layer): **1.31**

Disturbance severity: **0.7**

Disturbance severity (herb layer): **0.14**

Mowing frequency: **0.01**

Grazing pressure: **0.19**

Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**CA \(FAG\) *Carpino-Fagetea sylvaticae***](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**T1G *Alnus cordata* forest**](#)

Constant species of EUNIS habitats: [**T1G *Alnus cordata* forest**](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Synanthropic**

Distribution

Continentality: **3**

Continentality amplitude: **3**