Viola reichenbachiana

Habitus and growth type

Plant height [m]: **0.16**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **29.79**

Flower

Flowering period: **February-July**

Fruit, seed and dispersal

Seed mass [mg]: **2.27**
Dispersal mode: **Myrmecochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Moisture indicator value: **5.3**
Reaction indicator value: **7.1**
Nutrient indicator value: **5.6**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.03**
Disturbance frequency (herb layer): **1.31**
Disturbance severity: **0.69**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.13**
Mowing frequency: **0.01**
Grazing pressure: **0.2**
Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CA (FAG) *Carpino-Fagetea sylvaticae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T17 Fagus forest on non-acid soils**, **T1E Carpinus and Quercus mesic deciduous forest**, **T1F Ravine forest**, **T1G Alnus cordata forest**, **T32 Temperate mountain Abies forest**, **T3C Taxus baccata forest**
Constant species of EUNIS habitats: **S37 Corylus avellana scrub**, **T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils**, **T13 Temperate hardwood riparian forest**, **T17 Fagus forest on non-acid soils**, **T19 Temperate and submediterranean thermophilous deciduous forest**, **T1D Southern European mountain Betula and Populus tremula forest on mineral soils**, **T1E Carpinus and Quercus mesic deciduous forest**, **T1F Ravine forest**, **T1G Alnus cordata forest**, **T31 Temperate mountain Picea forest**, **T32 Temperate mountain Abies forest**, **T33 Mediterranean mountain Abies forest**, **T39 Mediterranean and Balkan subalpine Pinus heldreichii-Pinus peuce forest**, **T3C Taxus baccata forest**

Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Synanthropic**

Distribution

Continentality: **4**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz