Ziziphora capitata

Habitus and growth type

Plant height [m]: **0.15**
Life span: **Annual**
Life form: **Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **21.31**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.5**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**
Temperature indicator value: **7.4**
Moisture indicator value: **3.3**
Reaction indicator value: **6**
Nutrient indicator value: **2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.82**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.93**
Disturbance severity: **0.68**
Disturbance severity (herb layer): **0.58**
Mowing frequency: **0.42**
Grazing pressure: **0.33**
Soil disturbance: **0.32**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **T38 Mediterranean montane Cedrus forest**

Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz