

Crepis caucasica

Habitus and growth type

Plant height [m]: **0.54**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]:

Fruit, seed and dispersal

Seed mass [mg]: **0.65**

Dispersal mode: **Anemochory**

Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.9**

Temperature indicator value: **4**

Moisture indicator value: **7.9**

Reaction indicator value: **6.3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.99**

Disturbance frequency (herb layer): **1.38**

Disturbance severity: **0.29**

Disturbance severity (herb layer): **0.21**

Mowing frequency: **0.17**

Grazing pressure: **0.16**

Soil disturbance: **0.06**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**CM \(MOL\) Molinio-Arrhenatheretea**](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils**](#)

Constant species of EUNIS habitats: [**P2N Spring, Q41 Alkaline, calcareous, carbonate-rich small-sedge spring fen, Q42 Extremely rich moss-sedge fen, R35 Moist or wet mesotrophic to eutrophic hay meadow, T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils**](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Wetland, Mire, Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Forest**

Distribution

Continentality: **5**

Continentality amplitude: **4**