

# *Gnaphalium caucasicum*

## Habitus and growth type

Plant height [m]: **0.38**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **23.91**

## Fruit, seed and dispersal

Seed mass [mg]: **0.05**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.5**

Temperature indicator value: **4.7**

Reaction indicator value: **3.8**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.66**

Disturbance frequency (herb layer): **1.66**

Disturbance severity: **0.57**

Disturbance severity (herb layer): **0.42**

Mowing frequency: **0.36**

Grazing pressure: **0.25**

Soil disturbance: **0.27**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [QG \(EPI\) \*Epilobietea angustifolii\*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Heathland, Scrub, Forest, Synanthropic**

## **Distribution**

Continentality amplitude: **6**