Helianthus subcanescens

Habitus and growth type

Plant height [m]: **1.67**
Life span: **Perennial**
Life form: **Geophyte**

Leaf

Specific leaf area [mm^2/mg]: **32.81**

Fruit, seed and dispersal

Seed mass [mg]: **3.8**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Neophyte**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**
Temperature indicator value: **6.9**
Moisture indicator value: **6.4**
Reaction indicator value: **7.2**
Nutrient indicator value: **7.7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.04**
Disturbance frequency (herb layer): **1.52**
Disturbance severity: **0.67**
Disturbance severity (herb layer): **0.61**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.55**
Grazing pressure: **0.2**
Soil disturbance: **0.27**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QG (EPI) *Epilobietea angustifolii***

EUNIS habitat

Diagnostic species of EUNIS habitats: **V39 Mesic perennial anthropogenic herbaceous vegetation**

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: **5**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz