

Zannichellia clausii

Habitus and growth type

Plant height [m]: **0.3**

Life span: **Annual, Perennial**

Life form: **Hydrophyte**

Leaf

Specific leaf area [mm²/mg]: **81.13**

Fruit, seed and dispersal

Dispersal mode: **Hydrochory**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Saline**

Ellenberg-type indicator values

Light indicator value: **7.5**

Moisture indicator value: **11.6**

Reaction indicator value: **8**

Nutrient indicator value: **6**

Salinity indicator value: **8**

Disturbance indicator values

Disturbance frequency: **1.56**

Disturbance frequency (herb layer): **1.56**

Disturbance severity: **0.35**

Disturbance severity (herb layer): **0.32**

Mowing frequency: **0**

Grazing pressure: **0.03**

Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [MC \(RUP\) *Ruppiaetea maritimae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [P3a Brackish-water vegetation](#), [V14 Inundated or inundatable cropland, including rice fields](#)

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Aquatic, Saline vegetation, Synanthropic**