Asplenium adiantum-nigrum subsp. *serpentini*

Habitus and growth type

Plant height [m]: **0.23**

Flower

Flowering period: **May-September**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **3**

Habitat and sociology

Syntaxon
Diagnostic species of phytosociological classes: **CO (ERI) *Erico-Pinetea***
EUNIS habitat
Diagnostic species of EUNIS habitats: **R17 Heavy-metal dry grassland of the Balkans**, **U3A Temperate ultramafic inland cliff**, **U3B Mediterranean ultramafic inland cliff**
Constant species of EUNIS habitats: **R17 Heavy-metal dry grassland of the Balkans**, **U3A Temperate ultramafic inland cliff**, **U3B Mediterranean ultramafic inland cliff**
Dominant species of EUNIS habitats: **U3A Temperate ultramafic inland cliff**
Broad habitat
Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **4**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz