Atropa bella-donna subsp. *caucasica*

Habitus and growth type

Plant height [m]: **1.05**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **28.13**

Fruit, seed and dispersal

Seed mass [mg]: **493.03**
Dispersal mode: **Endozoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.9**
Temperature indicator value: **5.6**
Moisture indicator value: **5.2**
Reaction indicator value: **7.4**
Nutrient indicator value: **7.6**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.11**
Disturbance frequency (herb layer): **1.32**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.22**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.07**
Grazing pressure: **0.19**
Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QG (EPI) *Epilobietea angustifolii***

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest, Synanthropic**

Distribution

Continentality: **5**
Continentality amplitude: **3**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz