Bromus lanceolatus subsp. *lanceolatus*

Habitus and growth type

Plant height [m]: **0.5**
Life span: **Annual**
Life form: **Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **26.19**

Fruit, seed and dispersal

Seed mass [mg]: **3.93**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**
Temperature indicator value: **9**
Moisture indicator value: **3**
Reaction indicator value: **7**
Nutrient indicator value: **5.5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.83**
Disturbance frequency (herb layer): **1.89**
Disturbance severity: **0.76**
Disturbance severity (herb layer): **0.74**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **1**
Grazing pressure: **0.27**
Soil disturbance: **0.33**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QC (CHE) *Chenopodietea***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz