

Corynephorus articulatus subsp. *articulatus*

Habitus and growth type

Plant height [m]: **0.26**

Life span: **Annual**

Life form: **Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **10.5**

Moisture indicator value: **1.5**

Reaction indicator value: **8**

Nutrient indicator value: **1.5**

Salinity indicator value: **1**

Disturbance indicator values

Disturbance frequency: **1.27**

Disturbance frequency (herb layer): **2.03**

Disturbance severity: **0.56**

Disturbance severity (herb layer): **0.51**

Mowing frequency: **0.31**

Grazing pressure: **0.3**

Soil disturbance: **0.36**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FI \(TUB\) *Helianthemetea guttati*](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Scrub**