

Dianthus viscidus subsp. *viscidus*

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Disturbance indicator values

Disturbance frequency: **0.77**

Disturbance frequency (herb layer): **1.86**

Disturbance severity: **0.41**

Disturbance severity (herb layer): **0.31**

Mowing frequency: **0.03**

Grazing pressure: **0.33**

Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FO \(ANA\) *Trifolio anatolici-Polygonetea arenastri*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1K Balkan and Anatolian oromediterranean dry grassland](#)

Constant species of EUNIS habitats: [R1K Balkan and Anatolian oromediterranean dry grassland](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland**