Eryngium viviparum subsp. *bariegoi*

Habitus and growth type

Life span: **Biennial or short-lived**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Epizoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **6**
Moisture indicator value: **9**
Reaction indicator value: **5**
Nutrient indicator value: **2**
Salinity indicator value: **0**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **OC (ISO) *Isoëto-Nanojuncetea***

Broad habitat

Occurrence in broad habitats: **Wetland**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz