Halacsya sendtneri subsp. *devollensis*

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **7.9**
Temperature indicator value: **6.1**
Moisture indicator value: **4.4**

Disturbance indicator values

Disturbance frequency: **1.69**
Disturbance frequency (herb layer): **1.69**
Disturbance severity: **0.35**
Disturbance severity (herb layer): **0.35**
Mowing frequency: **0.42**
Grazing pressure: **0.31**
Soil disturbance: **0.27**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R17 Heavy-metal dry grassland of the Balkans**
Constant species of EUNIS habitats: **R17 Heavy-metal dry grassland of the Balkans**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz