Hypochaeris maculata subsp. *pelivanovicii*

Habitus and growth type

Plant height [m]: **0.38**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **20.65**

Fruit, seed and dispersal

Seed mass [mg]: **1.55**
Dispersal mode: **Myrmecochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.7**
Temperature indicator value: **5.8**
Moisture indicator value: **4.2**
Reaction indicator value: **6.8**
Nutrient indicator value: **2.5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.8**
Disturbance frequency (herb layer): **1.79**
Disturbance severity: **0.52**
Disturbance severity (herb layer): **0.42**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.59**
Grazing pressure: **0.28**
Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CI (NAR) *Nardetea strictae***, **DA (FES) *Festuco-Brometea***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

Distribution

Continentality: **6**
Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz