Chrysopogon gryllus subsp. *gryllus*

Habitus and growth type

Plant height [m]: **0.85**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **18.82**

Fruit, seed and dispersal

Seed mass [mg]:
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**
Moisture indicator value: **3.8**
Reaction indicator value: **7.3**
Nutrient indicator value: **3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.13**
Disturbance frequency (herb layer): **1.83**
Disturbance severity: **0.51**
Disturbance severity (herb layer): **0.47**
Mowing frequency: **0.63**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.32**
Soil disturbance: **0.17**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region**, **S34 Balkan-Anatolian submontane genistoid scrub**
Constant species of EUNIS habitats: **R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region**, **S34 Balkan-Anatolian submontane genistoid scrub**
Dominant species of EUNIS habitats: **R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz