

# *Liparis loeselii* subsp. *nemoralis*

## Habitus and growth type

Plant height [m]: **0.1**

Life span: **Perennial**

Life form: **Geophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **37.99**

## Fruit, seed and dispersal

Seed mass [mg]: **0**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **initial or partial mycoheterotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.9**

Temperature indicator value: **5.6**

Moisture indicator value: **8.7**

Reaction indicator value: **8.3**

Nutrient indicator value: **2.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.45**

Disturbance frequency (herb layer): **1.71**

Disturbance severity: **0.28**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.33**

Grazing pressure: **0.17**

Soil disturbance: **0.14**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [PA \(SCH\) \*Scheuchzerio palustris\*-\*Caricetea fuscae\*](#)

Broad habitat

Occurrence in broad habitats: **Spring, Mire**

## Distribution

Continentality: **5**

Continentality amplitude: **4**