Polypogon maritimus subsp. *maritimus*

Habitus and growth type

Plant height [m]: **0.21**
Life span: **Annual**
Life form: **Therophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.09**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Saline**

Ellenberg-type indicator values

Light indicator value: **8.3**
Moisture indicator value: **6**
Reaction indicator value: **7**
Nutrient indicator value: **5.8**
Salinity indicator value: **4.5**

Disturbance indicator values

Disturbance frequency: **1.68**
Disturbance frequency (herb layer): **1.75**
Disturbance severity: **0.47**
Disturbance severity (herb layer): **0.46**
Mowing frequency: **0.14**
Grazing pressure: **0.23**
Soil disturbance: **0.23**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **JA (SAG) *Saginetea maritimae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **MA251 Mediterranean upper saltmarsh**, **Q63 Periodically exposed saline shore with pioneer or ephemeral vegetation**

Constant species of EUNIS habitats: **MA251 Mediterranean upper saltmarsh**, **Q63 Periodically exposed saline shore with pioneer or ephemeral vegetation**

Dominant species of EUNIS habitats: **Q63 Periodically exposed saline shore with pioneer or ephemeral vegetation**

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Wetland, Saline vegetation**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz