Polypogon viridis subsp. *pauciflorus*

Habitus and growth type

Plant height [m]: **0.5**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.09**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**
Moisture indicator value: **7**
Salinity indicator value: **0.8**

Disturbance indicator values

Disturbance frequency: **1.17**
Disturbance frequency (herb layer): **1.65**
Disturbance severity: **0.45**
Disturbance severity (herb layer): **0.41**
Mowing frequency: **0.25**
Grazing pressure: **0.17**
Soil disturbance: **0.21**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CM (MOL) *Molinio-Arrhenatheretea***

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

EUNIS habitat

Diagnostic species of EUNIS habitats: **T25 Phoenix theophrasti vegetation**

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz