

Sedum magellense subsp. *olympicum*

Habitus and growth type

Plant height [m]: **0.13**

Life span: **Perennial**

Life form: **Chamaephyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.5**

Temperature indicator value: **5.5**

Moisture indicator value: **3**

Reaction indicator value: **7.5**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.5**

Disturbance frequency (herb layer): **1.35**

Disturbance severity: **0.43**

Disturbance severity (herb layer): **0.26**

Mowing frequency: **0.02**

Grazing pressure: **0.19**

Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [KC \(ASP\) *Asplenietea trichomanis*](#),
[KE \(THL\) *Thlaspietea rotundifolii*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [U29 Eastern Mediterranean base-rich scree](#)

Constant species of EUNIS habitats: [U29 Eastern Mediterranean base-rich scree](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Sparsely vegetated (incl. rock and scree)**