

# *Schoenoplectus lacustris* subsp. *glaucus*

## Habitus and growth type

Plant height [m]: **1.18**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **5.42**

## Flower

Flowering period: **April-August**

## Fruit, seed and dispersal

Seed mass [mg]: **1.54**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Ecology

Disturbance indicator values

Disturbance frequency: **1.83**

Disturbance frequency (herb layer): **1.9**

Disturbance severity: **0.34**

Disturbance severity (herb layer): **0.34**

Mowing frequency: **0.45**

Grazing pressure: **0.25**

Soil disturbance: **0.14**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [OD \(PHR\) Phragmito-Magnocaricetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [Q46 Carpathian travertine fen with halophytes](#), [Q54 Inland saline or brackish helophyte bed](#)

Constant species of EUNIS habitats: [MA232 Baltic coastal meadow](#), [Q46 Carpathian travertine fen with halophytes](#), [Q54 Inland saline or brackish helophyte bed](#)

Dominant species of EUNIS habitats: [Q46 Carpathian travertine fen with halophytes](#), [Q54 Inland saline or brackish helophyte bed](#)

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Aquatic, Wetland, Mire, Saline vegetation**

## **Distribution**

Continentality: 7

Continentality amplitude: **8**