

Verbascum lanatum subsp. *hinkei*

Habitus and growth type

Plant height [m]: **0.8**

Life span: **Biennial or short-lived, Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **5**

Moisture indicator value: **3**

Reaction indicator value: **5**

Nutrient indicator value: **7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.18**

Disturbance frequency (herb layer): **1.24**

Disturbance severity: **0.61**

Disturbance severity (herb layer): **0.21**

Mowing frequency: **0.07**

Grazing pressure: **0.21**

Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QG \(EPI\) Epilobietea angustifolii](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: **6**

Continentality amplitude: **2**