Polygonatum odoratum var. *odoratum*

Habitus and growth type

Plant height [m]: **0.36**
Life span: **Perennial**
Life form: **Geophyte**

Leaf

Specific leaf area [mm^2/mg]: **33.11**

Fruit, seed and dispersal

Seed mass [mg]: **309.22**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.5**
Moisture indicator value: **3.7**
Reaction indicator value: **7.3**
Nutrient indicator value: **3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.15**
Disturbance frequency (herb layer): **1.37**
Disturbance severity: **0.63**
Disturbance severity (herb layer): **0.18**
Mowing frequency: **0.06**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.22**
Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **BC (BRA) *Brachypodio pinnati-Betuletea pendulae***, **CC (QUE) *Quercetea robori-petraeae***, **CL (GER) *Trifolio-Geranietea sanguinei***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R51 Thermophilous forest fringe of base-rich soils**, **S24 Subalpine genistoid scrub of the Amphi-Adriatic region**, **T36 Temperate and submediterranean montane Pinus sylvestris-Pinus nigra forest**
Constant species of EUNIS habitats: **R51 Thermophilous forest fringe of base-rich soils**, **S24 Subalpine genistoid scrub of the Amphi-Adriatic region**, **T19 Temperate and submediterranean thermophilous deciduous forest**, **T36 Temperate and submediterranean montane Pinus sylvestris-Pinus nigra forest**, **T3H Larix light taiga**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality: **6**
Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz