

# *Trifolium stellatum* var. *xanthinum*

## Habitus and growth type

Plant height [m]: **0.15**

Life span: **Annual**

Life form: **Therophyte**

## Fruit, seed and dispersal

Seed mass [mg]: **3.07**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **9**

Moisture indicator value: **3.5**

Reaction indicator value: **6**

Nutrient indicator value: **5.5**

Salinity indicator value: **0.5**

Disturbance indicator values

Disturbance frequency: **0.77**

Disturbance frequency (herb layer): **1.83**

Disturbance severity: **0.75**

Disturbance severity (herb layer): **0.59**

Mowing frequency: **0.28**

Grazing pressure: **0.31**

Soil disturbance: **0.32**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FI \(TUB\) \*Helianthemetea guttati\*](#), [FJ \(TRA\) \*Stipo-Trachynietea distachyae\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [S72 Eastern Mediterranean spiny heath \(phrygana\)](#)

Constant species of EUNIS habitats: [R1F Mediterranean annual-rich dry grassland](#), [S63 Eastern garrigue](#), [S72 Eastern Mediterranean spiny heath \(phrygana\)](#), [T25 \*Phoenix theophrasti\* vegetation](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**