

# *Vicia grandiflora* var. *grandiflora*

## Habitus and growth type

Plant height [m]: **0.48**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **27.95**

## Fruit, seed and dispersal

Seed mass [mg]: **18.62**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.1**

Moisture indicator value: **4.1**

Reaction indicator value: **5.3**

Nutrient indicator value: **4.3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.71**

Disturbance frequency (herb layer): **1.77**

Disturbance severity: **0.65**

Disturbance severity (herb layer): **0.53**

Mowing frequency: **0.46**

Grazing pressure: **0.18**

Soil disturbance: **0.41**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [CB \(PUB\) Quercetea pubescentis](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Synanthropic**

## **Distribution**

Continentality: **7**

Continentality amplitude: **4**