# *Vicia melanops* var. *melanops*

### Habitus and growth type

Plant height [m]: **0.39**
Life span: **Annual**
Life form: **Therophyte**

### Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

### Taxon origin

Origin in Europe: **Native**

### Ecology

#### Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

#### Ellenberg-type indicator values

Light indicator value: **7**
Temperature indicator value: **8**
Moisture indicator value: **3**
Reaction indicator value: **5**
Nutrient indicator value: **5**
Salinity indicator value: **0**

#### Disturbance indicator values

Disturbance frequency: **0.35**
Disturbance frequency (herb layer): **1.36**
Disturbance severity: **0.67**
Disturbance severity (herb layer): **0.4**
Mowing frequency: **0.21**
Grazing pressure: **0.17**
Soil disturbance: **0.3**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **CL (GER) *Trifolio-Geranietea sanguinei***

### Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*