

# *Anogramma*

## **Habitus and growth type**

Plant height [m]: **0.14**

Life span: **Annual**

Life form: **Therophyte**

## **Flower**

Flowering period: **February-July**

## **Fruit, seed and dispersal**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **Native**

## **Ecology**

### Environmental relationships

Substrate humidity relationship: **Dry**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Moisture indicator value: **3**

Nutrient indicator value: **2.8**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.49**

Disturbance frequency (herb layer): **1.6**

Disturbance severity: **0.72**

Disturbance severity (herb layer): **0.51**

Mowing frequency: **0.18**

Grazing pressure: **0.28**

Soil disturbance: **0.26**

## **Habitat and sociology**

## Syntaxon

Diagnostic species of phytosociological classes: [KB \(POD\) Polypodietea](#)

## EUNIS habitat

Diagnostic species of EUNIS habitats: [T25 Phoenix theophrasti vegetation](#)

## Broad habitat

Occurrence in broad habitats: **Forest, Sparsely vegetated (incl. rock and scree)**