Calepina

Habitus and growth type

Plant height [m]: **0.29**
Life span: **Annual, Biennial or short-lived**
Life form: **Hemicryptophyte, Therophyte**

Flower

Flowering period: **January-May**

Fruit, seed and dispersal

Seed mass [mg]: **4.39**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8**
Moisture indicator value: **3.3**
Reaction indicator value: **7.3**
Salinity indicator value: **0.6**

Disturbance indicator values

Disturbance frequency: **1.59**
Disturbance frequency (herb layer): **1.94**
Disturbance severity: **0.77**
Disturbance severity (herb layer): **0.76**
Mowing frequency: **0.6**
Grazing pressure: **0.21**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Soil disturbance: **0.57**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QF (ART) *Artemisietea vulgaris***

Broad habitat

Occurrence in broad habitats: **Wetland, Synanthropic**

Distribution

Continentality: **6**
Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz