Cladium

Habitus and growth type

Plant height [m]: **1.4**
Life span: **Perennial**
Life form: **Hemicryptophyte, Geophyte**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **7.46**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **2.19**
Dispersal mode: **Myrmecochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.6**
Moisture indicator value: **9.6**
Reaction indicator value: **8.6**
Nutrient indicator value: **3.8**
Salinity indicator value: **0.9**

Disturbance indicator values

Disturbance frequency: **1.19**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.65**
Disturbance severity: **0.26**
Disturbance severity (herb layer): **0.22**
Mowing frequency: **0.46**
Grazing pressure: **0.12**
Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **OD (PHR) *Phragmito-Magnocaricetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **N1J Mediterranean and Black Sea moist and wet dune slack**, **Q43 Tall-sedge base-rich fen**, **Q53 Tall-sedge bed**
Constant species of EUNIS habitats: **Q43 Tall-sedge base-rich fen**
Dominant species of EUNIS habitats: **N1J Mediterranean and Black Sea moist and wet dune slack**, **Q53 Tall-sedge bed**

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Mire**

Distribution

Continentality amplitude: **7**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz